



Infection Prevention for Drive-Thru PODs

Vaccination Staff	
Staff who <u>will</u> have contact with asymptomatic* individuals	
Facemask?	Yes for extended use unless visibly soiled or contaminated
Gloves?	Yes, and changed between each encounter. Perform hand hygiene immediately after removing gloves
Gown?	No
Eye Protection?	Goggles or Face Shields for extended use unless visibly soiled or contaminated
Hand Hygiene?	60-95% alcohol
Physical Distancing?	Maintain 6 feet until necessary to provide vaccine or medical care
POD Staff	
Staff who <u>may</u> have contact with asymptomatic* individuals	
Facemask?	Yes for extended use unless visibly soiled or contaminated
Gloves?	When contact with patients, specimens, or potentially contaminated surfaces is expected.
Gown?	No
Eye protection?	Goggles or Face Shields for extended use unless visibly soiled or contaminated
Hand Hygiene?	60-95% alcohol
Physical Distancing?	Maintain 6 feet



ENVIRONMENT	
Location?	Outside, Vehicles, Pop-up Tents
Surfaces?	Disinfected at least two times a shift
Shared supplies?	Disinfected after each use is recommended

*Generally, routine vaccination should be deferred for individuals with suspect or known COVID-19 until [criteria](#) have been met. Persons who have had a known exposure to a person with COVID-19 should not seek outpatient care solely for vaccination until their 14-day quarantine period has ended. Additional considerations available [here](#).

DETAILS:
<p><u>Facemask:</u> Facemasks are worn consistently for extended timeframes. When the facemask is visibly soiled or contaminated, doff and don new facemask immediately. Perform hand hygiene before and after mask readjustment or removal.</p>
<p><u>Glove:</u> Gloves are worn consistently when providing care or when contact with contaminated surfaces or specimens is expected. Hand hygiene should be performed immediately after removal of gloves. Gloves should be doffed after each patient encounter or after contact with a contaminated surface.</p>
<p><u>Eye protection:</u> Goggles and face shield are worn consistently for extended timeframe. If eye protection becomes visibly soiled or contaminated, disinfect or change eye protection immediately. Perform hand hygiene before and after eye protection readjustment or removal.</p>
<p><u>Extended use:</u> refers to the practice of wearing the same mask and eye protection for repeated close contact encounters with several patients, without removing the respirator between patient encounters. Extended use PPE should be removed when visibly contaminated, integrity is disrupted, or after contact with a symptomatic or COVID-19 positive individual in outpatient scenarios.</p>
<p><u>Surfaces:</u> At a minimum, surfaces commonly touched are disinfected twice during a shift and as needed when contamination occurs. It is preferred to have all surfaces wiped down after use or as soon as possible to decrease transmission of viruses.</p>



Shared supplies: It is recommended shared supplies and equipment are disinfected after each use. Shared supplies include pens and clipboards. PPE should never be shared between individuals.

Standard Precautions: the basic level of infection control precautions that should be used in the care of all patients at all times. Applies to blood, all body fluids, secretions, and excretions (except sweat), whether or not they contain visible blood; non-intact skin and mucous membranes.

Personal protective equipment (PPE) Medical-grade supplies, including gloves, gowns, masks, and eye protection use to adhere to standard precautions and universal PPE guidance.

For more information see Infection Control Guidance at
<https://www.oregon.gov/oha/covid19/Pages/Healthcare-Partners.aspx>

Document accessibility: For individuals with disabilities or individuals who speak a language other than English, OHA can provide information in alternate formats such as translations, large print, or braille. Contact the Health Information Center at 1-971-673-2411, 711 TTY or COVID19.LanguageAccess@dhs.oha.state.or.us