

COVID-19 Vaccine Doses

COVID-19 Vaccine	First dose	Second dose	Third dose	Additional dose ¹ (for immunocompromised only)	Booster doses ²
Moderna monovalent pediatric primary series (6 months–5 years)	0.25 mL (Dark blue cap, magenta label border)	0.25 mL	N/A	0.25 mL	N/A
Pfizer monovalent pediatric primary series (6 months–4 years)	0.2 mL (Maroon cap)	0.2 mL	0.2 mL	N/A	N/A
Pfizer pediatric primary series (ages 5–11)	0.2 mL (Orange cap)	0.2 mL	N/A	0.2 mL	0.2 mL
Moderna monovalent pediatric primary series (ages 6–11)	0.5 mL (dark blue cap, teal or purple label border)	0.5 mL	N/A	0.5 mL	N/A
Pfizer bivalent booster dose (ages 5–11)	N/A	N/A	N/A	N/A	0.2 mL (orange cap, orange label border)
Moderna bivalent booster dose (6–11)	N/A	N/A	N/A	N/A	0.5 mL (dark blue cap, gray label border)
Pfizer monovalent primary dose (ages 12+)	0.3 mL (Gray cap)	0.3 mL	N/A	0.3 mL	This formulation not authorized as a booster dose; see bivalent booster options below
Moderna monovalent primary dose (ages 12+)	0.5 mL (Red cap, light blue label border)	0.5 mL	N/A	0.5 mL	This formulation not authorized as a booster dose; see bivalent booster options below.
Pfizer bivalent booster dose (ages 12+)	N/A	N/A	N/A	N/A	0.3 mL (Gray cap)

Moderna bivalent booster dose (ages 12+)	N/A	N/A	N/A	N/A	0.5 mL (dark blue cap, gray label border)
Johnson & Johnson (ages 18+)	0.5 mL	N/A	N/A	See note below	See note below
Novavax³ (ages 12+)	0.5 mL	0.5 mL	N/A	N/A	N/A

Note: Immunocompromised people who received the Johnson & Johnson vaccine should get an additional dose of a monovalent (original vaccine formula) mRNA vaccine (Pfizer or Moderna) at least 28 days after their initial dose. Then a booster dose of a bivalent (updated vaccine formula) mRNA vaccine at least two months after their additional dose OR at least two months after their most recent monovalent booster dose.

¹ CDC recommends that people who are immunocompromised and received Pfizer (5+) or Moderna (6 months+) get an additional (monovalent) dose at least 28 days after their second dose. Additional doses for immunocompromised persons are part of the primary vaccine series. Additional doses of Pfizer vaccine for children 6 months through 4 years of age are not recommended at this time.

² CDC recommends that everyone age 5 and older who received Pfizer or Moderna, and 18 and over who received Johnson & Johnson, get a bivalent Moderna or Pfizer booster dose when they are eligible—i.e., at least 2 months after completion of the primary series (including additional doses as indicated for immunocompromised persons) or 2 months after receipt of the most recent monovalent booster dose.

³ Additional doses for immunocompromised persons are not currently recommended for Novavax vaccine recipients.

Document accessibility: For individuals with disabilities or individuals who speak a language other than English, OHA can provide information in alternate formats such as translations, large print, or braille. Contact the COVID-19 Communications Unit at 971-673-2411, 711 TTY or COVID19.LanguageAccess@dhs.ohio.gov