



## COVID-19 Vaccine Doses

COVID-19 Vaccine	First dose	Second dose	Third dose* (for immunocompromised only)	Booster dose*
<b>Pfizer pediatric (Ages 5–11)</b>	0.2 mL	0.2 mL	0.2 mL	Not recommended by FDA or CDC
<b>Pfizer (Ages 12+)</b>	0.3 mL	0.3 mL	0.3 mL	0.3 mL (for people ages 12+, five months after second dose)
<b>Moderna (Ages 18+)</b>	0.5 mL	0.5 mL	0.5 mL	0.25 mL (for people ages 18+, five months after second dose)
<b>Janssen (J&amp;J; Ages 18+)</b>	0.5 mL		Not recommended by FDA or CDC	0.5 mL (for people ages 18+, two months after first dose)

\*The CDC recommends everyone age 12 and older get a booster dose when they are eligible: five months after their second dose of a Pfizer or Moderna vaccine, or two months after a Janssen (Johnson & Johnson) vaccine dose. This includes people who are immunocompromised and received a third dose of Pfizer or Moderna.

**Document accessibility:** For individuals with disabilities or individuals who speak a language other than English, OHA can provide information in alternate formats such as translations, large print, or braille. Contact the COVID-19 Communications Unit at 1-971-673-2411, 711 TTY or [COVID19.LanguageAccess@dhsosha.state.or.us](mailto:COVID19.LanguageAccess@dhsosha.state.or.us).